


## Nanangenine F

Code No.: **BIA-N1969**

Pack sizes: **0.5 mg, 2.5 mg**



Synonyms :

### Specifications

CAS #	:	<b>NA</b>
Molecular Formula	:	<b>C<sub>21</sub>H<sub>34</sub>O<sub>5</sub></b>
Molecular Weight	:	<b>366.5</b>
Source	:	<b><i>Aspergillus nanangensis</i></b>
Appearance	:	<b>Off-white solid</b>
Purity	:	<b>&gt;95% by HPLC</b>
Long Term Storage	:	<b>-20°C</b>
Solubility	:	<b>Soluble in methanol or DMSO</b>

### Application Notes

The nanangenines are newly discovered drimane sesquiterpenoid metabolites from a hitherto undescribed Australian fungus, *Aspergillus nanangensis*. Nanangenine F is an acylated drimane and putative biosynthetic intermediate of the higher nanangenines. Nanangenine F is weakly active against *B. subtilis* (IC<sub>50</sub> 78 µg/mL) and tumor cell lines (mouse myeloma NS-1 cell, human prostate DU-145 and human breast adenocarcinoma MC-7; IC<sub>50</sub> 49 - 95 µg/mL) and neonatal foreskin cell line. The nanangenines did not exhibit Gram negative antibacterial, antifungal or herbicidal activity.

### References

1. Nanangenines: drimane sesquiterpenoids as the dominant metabolite cohort of a novel Australian fungus, *Aspergillus nanangensis*. Lacey H.J. et al. *Beilstein J Org Chem*. 2019, 15, 2631.