


Pikromycin

Code No.: **BIA-P1915**

Pack sizes: **0.5 mg, 2.5 mg**



Synonyms : Albomycetin, Amaromycin, Antibiotic B 62169A

Specifications

CAS #	: 19721-56-3
Molecular Formula	: C ₂₈ H ₄₇ NO ₈
Molecular Weight	: 525.7
Source	: <i>Streptomyces</i> sp.
Appearance	: White crystalline solid
Purity	: >95% by HPLC
Long Term Storage	: -20°C
Solubility	: Soluble in DMSO

Application Notes

Pikromycin was the first macrolide antibiotic, isolated in 1951 by the German researchers Brockmann and co-workers from *Streptomyces* sp.. Its structure as a 14-membered lactone linked to D-desosamine was confirmed in 1957. Pikromycin is synthesized through a type I polyketide synthase system in *Streptomyces venezuelae*. Pikromycin is active against Gram positive bacteria and Mycobacteria. Pikromycin inhibits translation in bacteria by binding to overlapping sites in the ribosomal exit tunnel. Although pikromycin readily arrests the growth of bacteria, ~40% of cellular proteins continue to be synthesized even at saturating concentrations. Pikromycin is a weak inhibitor of human prolyl endopeptidase.

References

1. Pikromycin, ein bitter schmeckendes Antibiotikum aus Actinomyceten (Antibiotica aus actinomyceten, VI. Mitteil. Brockmann H. & Henkel W. Chem. Ber. 1951, 84, 284.
2. Biosynthesis and combinatorial biosynthesis of pikromycin-related macrolides in *Streptomyces venezuelae*. Xue Y. & Sherman S. Metabolic Engineering 2001, 3, 15.
3. Co-produced natural ketolides methymycin and pikromycin inhibit bacterial growth by preventing synthesis of a limited number of proteins. Almutairi M.M. et al. Nucleic Acids Res. 2017, 45, 9573.
4. Albocycline- and carbomycin-type macrolides, inhibitors of human prolyl endopeptidases. Christner C. et al. J Antibiot. 1998, 51, 368.